

HOW POTENTIALLY AVOIDABLE EVENTS ARE CALCULATED

Emergent Care for Injury Caused by Fall *(as of 1/1/2017 this measure is no longer reported by CMS)

- 1. Checked for at transfer and discharge
- 2. Include if M2310 = 02 (fall)

Emergent Care for Wound Infections, Deteriorating Wound Status *(as of 1/1/2017 this measure is no longer reported by CMS)

- 1. Checked for at transfer and discharge
- 2. Include if M2310 = 15 (wound infection or deterioration)

Emergent Care for Improper Medication Administration, Medication Side Effects

- 1. Checked for at transfer and discharge
- 2. Include if M2310 = 01 (improper medication administration, side effects)

Emergent Care for Hypo/Hyperglycemia

- 1. Checked for at transfer and discharge
- 2. Include if M2310 = 10 (hypo/hyperglycemia, diabetes out of control)

Development of Urinary Tract Infection

- 1. Check at Discharge
- 2. Include if M1600 was 00 (no) on most recent SOC or ROC, but is 01 (yes) at discharge

Increase in Number of Pressure Ulcers

- 1. Compare responses at discharge to most recent SOC or ROC
- 2. For assessments 1/1/2017 or later. Include if total of lines M1311 lines a1, b1, c1, d1, e1, and f1 is higher at discharge
- 3. For assessments prior to 1/1/2017. Include if total of lines M1308 lines a, b, c, d1, d2, and d3 is higher at discharge

Substantial Decline in 3 or More Activities of Daily Living

- 1. Compare responses at discharge to most recent SOC or ROC
- 2. Include if there has been a 2 level or more decline in 3 or more of the following.
 - a. M1800 (grooming)
 - b. M1830 (bathing)
 - c. M1840 (toilet transferring)
 - d. M1845 (toilet hygiene)
 - e. M1850 (transferring)
 - f. M1860 (ambulation)

Substantial Decline in Management of Oral Medications

- 1. Compare responses at discharge to most recent SOC or ROC
- 2. Include if M2020 (management of oral meds) declined from 00 at SOC/ROC to 03 at discharge

Discharged to the Community Needing Wound Care or Medication Assistance

- 1. Check at Discharge
- 2. Include if:
 - a. M2420 (discharge to community) = 01 (patient remained in community) and
 - b. M2102 (type of assistance) line c (med admin) = 03 or 04 and
 - c. M2020 (management of oral meds) = 02 or 03 and
 - d. M1710(confusion) <u>is not</u> 03, 04, or NA
 - a. M2420 (discharge to community) = 01 (patient remained in community) and
 - b. M2102 (type of assistance) line d (medical procedures) = 03 or 04 and
 - c. M1710(confusion) is not 03, 04, or NA and
 - d. M1324 (stage of worst pressure ulcer) = 02, 03, or 04 or
 - e. M1342 (stage of worst surgical wound) = 02 or 03

Discharged to the Community Needing Toileting Assistance

- 1. Check at Discharge
- 2. Include if:
 - a. M2420 (discharge to community) = 01 (pt remained in community) and
 - b. M2102 line a (ADL assistance) = 03 or 04 and
 - c. M1840 (toilet transferring) = 03 or 04 or
 - d. M1845 (toilet hygiene) = 02 or 03

Discharged to the Community with Behavioral Problems

- 1. Check at Discharge
- 2. Include if:
 - a. M2420 (discharge to community) = 01 and
 - b. M2102 line f (supervision and assistance) = 03 or 04 and
 - c. More than one of the following selections from M1740 (cognition) = 01, 02, 03, 04, 05, or 06

Discharged to the Community with an Unhealed Stage II Pressure Ulcer

- 1. Check at Discharge
- 2. Include if:
 - a. M2420 (discharge to community) = 01 and
 - b. M1307 (oldest stage II pressure ulcer) = 01 and
 - Length of stay 30 days or more (M0030/M0032 as compared to M0906)
 - d. M1307 = 02 and recorded date is 30 days or greater prior to discharge.